


L 4355-66

ACC NR: AP5628784



The work was presented by P. Nikolov, Corresponding Member of BAN, 16 Oct 64. Orig. art. has: 2 tables, 1 figure. [JPRS]

SUB CODE: 1S / SUBM DATE: 16Oct64 / ORIG REF: 003 / OTH REF: 006

KC
Circ 2/2

MARKOV, M.; TOSHEV, Iv.

On postoperative evisceration. Khirurgiia (Sofia) 18 no.4:
459-465 '65

1. Okruzhn onkologichen dispanser, Varna (gl. lekar M. Markov).

DASKALOV, D.; MARKOV, M.; MINEV, Ts.

Neurovegetative reaction in women and men. Izv biol med BAN 3 no.4:
91-97 '60. (EEAI 10:3)

1. Institut po fiziologiya pri BAN (Direktor: akad. D.Orakhovets)
(NERVOUS SYSTEM)

MARKOV, M.

Changes in the electroconductivity of skin in contact with anod.
Doklady BAN 14 no.4:413-415 '61.

1. Physiologisches Institut an der Bulgarischen Akademie der
Wissenschaften. Vorgelegt von Akademiemitglied D. Orakhovats
[D. Orakhovats].

MARKOV, M.

Relative dependency of electroconductivity from number of active perspiratory glands after electrophoresis of acetylcholin adrenalin and pilocarpin. Doklady BAN 14 no.5:555-558 '61.

1. Physiologisches Institut der Bulgarischen Akademie der Wissenschaften. Vorgelegt von Akademiemitglied D. Orachovats[D. Orakhovats]

(Electric conductivity) (Cataphoresis)
(Adrenalin)


MARKOV, M.; PETROV, I.

Investigation of perspiration suggested during hypnotic condition.
Doklady BAN 14 no.7:771-774 '61.

1. Vorgelegt von Akademiemitglied D. Orakhovats [D. Orakhovats].

(PERSPIRATION) (HYPNOTISM)

MARKOV, M.



MARKOV, M.; PETROV, I.

On the mechanism of gustatory sweating. Dokl. Bolg. akad.
nauk 15 no.1:89-92 '62.

1. Vorgelegt von Akademienmitglied D. Orahovats.
(SWEATING physiol) (TASTE physiol)

ORANOVATS, D.; MARKOV, M.; WARBANOV, W.

Effect of the resistance of the soft tissue of the extremities
in measuring blood pressure with Korotkoff's method. Dokl.
Bolg. akad. nauk 15 no.3:333-336 '62.

(BLOOD PRESSURE) (LEG blood supply)

MARKOV, M.

Changes in the electrical conductivity and potential of the skin in pressure on the anode in relation to the initial degree of electrical conductivity and potential differences. Izv. inst. fiziol. 5:179-186 '62.

(SKIN physiol)

MAR 24 1961

21. "The Nature of Psychological Control in the Main Adult Attachment Styles (L.A.)" ERIKSON article in Evolution, pp 61-80.
22. "On the Mechanics of 'Selfish' Feeling," ERIKSON and I. PIRELLA article in Evolution, pp 89-92.
23. "On the Nature of the Individual," ERIKSON article in Evolution, pp 93-95.
24. "Violence and the Subject," ERIKSON article in Evolution, pp 96-100.
25. "The Individual in the Social Context," ERIKSON article in Evolution, pp 101-105.

ORAKHOVATS, D. [Orakhovats, D.], akad.; MARKOV, M.; WARBANOV, W. [Vurbanov, V.]

Effect of the resistance of soft tissues in the measurement of blood pressure by the Korotkov method. Doklady BAN 15 no.3:333-336 '62.

1. Physiologisches Institut der Bulgarischen Akademie der Wissenschaften. 2. Chlen Redaktsionnoy kollegii, "Doklady Bolgarskoy Akademii Nauk" (for Orakhovats).

*

ORAKHOVATS, D. [deceased]; DRAGANOV, I.; VUKOVICH, V.; MARKOV, M.; FICOV, A.;
PAMENOV, V.; NACHEV, GE.

Simultaneous recording of the arterial pressure, splanchnic,
Korotkov's tones and pressure of the cuff of the brachial artery
by means of direct measurements in man. Izv. Inst. fiziol. Sofiya
7:5-18 '64.

ORAKHOVATS, D. [deceased]; MARKOV, M.; V'ERBANC, V.; DRAGANOV, L.

Studies on the development of Korotkov's tones. Izv. Inst. fiz. ...
(Sofia) 7:19-28 '64.

MARCHOV, M.

Large amounts of this substance are found in the
of active cells of glands following electrophoresis of various
concentrations of acetylcholine. (Int. Inst. Physiol. 1958, 13: 133-138).

ACC NR: AP7008901

SOURCE CODE: UR/0386/66/003/002/0072/0100

AUTHOR: Markov, M.

ORG: Physics Institute imeni P. N. Lebedev, Academy of Sciences USSR (Fizicheskii Institut AN SSSR)

TITLE: Search for specific mu-meson and sup nu mu-neutrino interactions at very high energies

SOURCE: Zhurnal eksperimental'noy i teoreticheskoy fiziki. Pis'ma v redaktsiya, v. 3, no. 2, 1966, 92-100

TOPIC TAGS: mu meson, meson interaction, neutrino, cosmic ray

SUB CODE: 21

ABSTRACT: New information gathered from cosmic ray studies seems to indicate that at very high energies there may exist specific Λ -meson and Σ -neutrino interactions in addition to the usual electromagnetic and weak interactions. The author investigates a pseudovector (pseudomaxwellian) field with $g(\pi/mc) \gamma_5 \gamma_\mu \gamma_\nu (\partial P_\mu / \partial x_\nu)$ -type interactions. In the first approximation the g -charge is carried by a Λ -meson and nucleons only; a Σ -neutrino and electron carry a much weaker g -charge. An evaluation of Λ -pair production by pseudophotons in Coulomb fields of extended nuclei is given for $Z = 10$, ($\sigma_{2\Lambda} \approx 10^{-27}$ - 10^{28} cm²), and a very general discussion of the existing experimental material is presented. Detailed numerical evaluations must be postponed until more reliable data becomes available. [JPRS: 35,630]

Card 1/1

MARKOV, M.; MININ, Ye.

Changed number of active sweat glands of the forearm in persons
of different ages after acetylcholine and adrenaline electroporation
before and after thermal stimulation. Izv. Inst. fiziol. (Sov. Union)
8:69-76 '64

RUSINOV, K.; NARKOV, V.; BASHKIN, IRAGANOVA, ... T SHKOVA, S.

Some pharmacophysiology of ...
(Soviet) ...

MARKOV, M.

Age changes in the electrical conductivity of the skin after the electrophoretic administration of acetylcholine, adrenaline and pilocarpine. Dokl. Bolg. akad. nauk 18 no.2:183-186 '65

1. Submitted on October 16, 1964

5(3) 5.5600

66965

AUTHORS: Kondrat'yev, D. A., Markov, M. A.,
Minachev, Kh. M.

SOV/32-25-11-13/69

TITLE: Analysis of Mixture of C₅ to C₇ Hydrocarbons by the Method of
Liquid - Gas Chromatography

PERIODICAL: Zavodskaya laboratoriya, 1959, Vol 25, Nr 11, pp 1301-1304
(USSR)

ABSTRACT: A simple device with a microflame detector (Ref 1) designed
for the analysis of paraffin hydrocarbons, naphthenes, aromatic,
and some unsaturated C₅ to C₇ hydrocarbons has been developed.
The construction of the dosing evaporator and the microflame
detector was based on a device developed by B. A. Rudenko
(Izvestiya AN SSSR, in the press). The detector is, in principle,
a burner consisting of a capillary 1 mm in diameter. The burner
is located at the upper output of a column 6 m long, and is
connected to a Chromel-Alumel thermocouple (Fig 1: Scheme of
the device). Data obtained with the detector are recorded by
automatic recording potentiometer of the types PS1-01^{7b} or
EPP-09^{7b} with a second PP potentiometer connected thereto. The
hydrocarbons may be analyzed by means of a thermostat of the

Card 1/3

66965

Analysis of Mixtures of C_5 to C_7 Hydrocarbons
by the Method of Liquid-Gas Chromatography

SOV/32-5-11-13/63

type TS-15²⁸ at constant or variable temperature. The best separating efficiency was reached when two columns (each 3 m long) were used, the one filled with diatomite brick chips (0.25 to 0.5mm) and tricresyl phosphate, and the other filled with diatomite brick chips and dioctyl phthalate. Separation was first effected at 15 - 20°C (for 15 minutes), and all normal and isoparaffin hydrocarbons C_5 to C_7 were separated from one another, whereafter temperature was raised to 85°C (1.5° per minute). Hydrogen was passed through the system with a rate of 60 cm³ per minute. The chromatogram of a 15-component (C_5 to C_7 hydrocarbon) mixture shows that all substances could be separated except for the pairs 2,3-dimethylbutane-2-methylpentane, cyclopentane-3-methylpentane, and cyclohexane-3-methylhexane. Results of an analysis of an artificial hydrocarbon mixture (Table 1) as well as with catalyzates at elevated temperature and hydrogen pressure (Table 2) are given. There are 3 figures, 2 tables, and 2 Soviet references. ✓

Card 2/3

Analysis of Mixtures of C₅ to C₇ Hydrocarbons
by the Method of Liquid - Gas Chromatography

66965

SOV/32-25-11-13/69

ASSOCIATION: Institut organicheskoy khimii im. N. D. Zelinskogo Akademii
nauk SSSR (Institute of Organic Chemistry imeni N.D.Zelinskiy
of the Academy of Sciences of the USSR) 4

Card 3/3

MINACHEV, Kh.M.; SHUYKIN, N.I.; MARKOV, M.A.

Effect of the specific surface of a platinized aluminosilicate on the degree of n-nonane conversion. Report No.1: Change in the activity of platinized aluminosilicate in the course of the treatment of the carrier with hydrogen. Izv.AN SSSR Otd.khim. nauk no.5:907-912 My '60. (MIRA 13:6)

1. Institut organicheskoy khimii imeni N.D. Zelinskogo Akademii nauk SSSR.

(Aluminosilicates) (Platinum) (Nonane)

S/062/60/000/008/023/033/XX
B013/B055

AUTHORS: Minachev Kh. M., Shuykin, N. I., and Markov, M. A.

TITLE: Investigation of the Effect of the Specific Surface of
Platinized Alumosilicate on the Degree of n-Nonane
Conversion

PERIODICAL: Izvestiya Akademii nauk SSSR. Otdeleniye khimicheskikh nauk.
1960. No. 8. pp. 1466-1470

TEXT: This paper is a continuation of the studies on the effect of the specific surface of catalysts on the degree of hydrocarbon conversion. For this purpose, the authors studied the conversion of n-nonane on alumosilicate catalysts with widely varying specific surfaces. Two 0.5% platinum/alumosilicate catalysts with specific surfaces of 320 m²/g (K-1) and 60 m²/g (K-2) were used. A standard alumosilicate catalyst for the cracking process was used as carrier for the preparation of the platinum catalysts. To prepare the latter, the carrier was saturated with a dilute H₂PtCl₆ solution. The data of the n-nonane used were in agreement with those given in Ref. 4. Infrared spectra showed the n-nonane to be free
Card 1/3

Investigation of the Effect of the Specific
Surface of Platinized Aluminosilicate on the
Degree of n-Nonane Conversion

S/062/60/000/008/023/033/XX
B013/B055

of isomers. The experimental apparatus has been described in Ref. 1. The experiments were carried out in a continuous system at 360 to 450°C. a hydrogen pressure of 10 atm and a flow rate of 1 h⁻¹. The molar ratio of hydrocarbon and hydrogen was 1:5. A fresh catalyst was used for each experiment. The results of the examination of the catalyzates are listed in Tables 1 and 2 and graphically represented in Figs. 1 - 3. These data show that the degree of hydrocracking of hydrocarbons considerably decreases with decreasing specific catalyst surface. This is in agreement with data given in Ref. 2. The yields of hydrocracking products on K-1 catalysts were found to increase more rapidly with a temperature rise than on K-2 catalysts. At temperatures of 420°C - 450°C, hydrocracking on K-2 is insignificant, which enables C₉ isoalkanes to be obtained in comparatively high yields (54% at 450°C). On K-1, the maximum yield of isononanes is obtained at 380°C (53%). Since aromatization occurs to a noticeable degree only at 400°C, aromatic hydrocarbons can be obtained over K-2 before C₉ isoalkane yields are reduced. The total yield of hydrocracking products under the experimental conditions was 45.7% over K-1, and 25.3% over K-2. The experiments have thus shown that by decreasing the specific surface

Card 2/3

Investigation of the Effect of the Specific S/062/60/000/008/023/033/XX
Surface of Platinized Aluminosilicate on the B013/B055
Degree of n-Nonane Conversion

of the catalyst the process can be carried out at higher temperatures without the occurrence of hydrocracking. There are 3 figures, 2 tables, and 4 Soviet references.

ASSOCIATION: Institut organicheskoy khimii im. N. D. Zelinskogo Akademii nauk SSSR
(Institute of Organic Chemistry imeni N. D. Zelinskii of the Academy of Sciences USSR;

SUBMITTED: February 18, 1959

Card 3/3

5. 34. 1

NOV 21 - 1964

AUTHORS: Tits-Skvortsova, I. N., Danilova, T. A., Markov, M. A.,
Stepanova, I. I., Osipenko, Ts. D.

TITLE: Synthesis and Conversions of Sulfur Compounds of Naphthalene Series Over an Alumina-Silica Catalyst

PERIODICAL: Zhurnal obshchey khimii, 1960, Vol 30, Nr 3, pp 985-991 (USSR)

ABSTRACT: The following compounds were synthesized and their conversions over an alumina-silica catalyst at 300° was studied. ***a*** - Thionaphtnol (72%), bp 143-144° (6 mm); ***β***-thionaphtnol (80%), mp 79-80°; ***α*** -naphthyl decyl sulfide (72%); ***α*** -naphthyl cyclopentyl sulfide (45.6%), bp 168-168.5° (2 mm), n_D^{20} 1.6419, d_4^{20} 1.1193;

β -naphthyl decyl sulfide (68%), bp 209-219° (2.5 mm), mp 34-35°; β -naphthyl cyclopentyl sulfide (65%), bp 187.5-188° (4 mm), n_D^{20} 1.6455, d_4^{20} 1.1052. This


Card 1/5

Synthesis and Conversions of Sulfur Compounds
of Naphthalene Series Over an Alumina-Silica
Catalyst


78297

SOV/79-30-3-51/69

study was undertaken to see whether the conversions of the thionaphthols over the above catalyst at 300° proceed similarly to the conversions of aromatic thiols under the same conditions. Conversions of aromatic thiols proceed as authors showed (DAN SSSR, 80, 377, 1951; ZhOKh, 21, 212, (1951); and others), according to the following scheme:



It was found that both α - and β -thionaphthols undergo an identical conversion over this catalyst at 300°, according to the following scheme:




Card 2/5

Synthesis and Conversions of Sulfur Compounds
of Naphthalene Series Over an Alumina-Silica
Catalyst

7
SOV, 1960-3-21/70

Comparison of schemes 1 and 2 shows that the isomeric α - and β -thionaphthols and aromatic thiols undergo similar conversions over the same catalyst at the same temperature. α -Naphthyl decyl sulfide decomposes over the catalyst at 300° to form naphthalene (36%, of weight of catalyst), decyl mercaptan (13.1%), decene (7.8%), and H₂S, according to scheme:



α -Naphthyl cyclopentyl sulfide decomposes over the catalyst to form naphthalene (40% of weight of catalyst), cyclopentanethiol (6.6%), dicyclopentyl sulfide (2.2%) and H₂S. The reaction proceeds also analogously to scheme 3. Catalytic decomposition of β -naphthyl cyclopentyl sulfide under above conditions results in the formation of β -thionaphthol (15.6% of weight of catalyst), cyclopentene (10.2%),


Card 3/5

Synthesis and Conversions of Sulfur Compounds
of Naphthalene Series Over an Alumina-Silica
Catalyst


78297

SOV/79-30-3-11/69

naphthalene (43.5%) and H_2S , according to a different
scheme:



Catalytic decomposition of β -naphthyl decyl sulfide
under the same conditions results in the formation of:
 β -thionaphthol (1.1% of weight of catalyst), decyl
mercaptan (6%), naphthalene (30.5%), decene-1 (4.2%)
fraction (4.2%) and H_2S , according to:




Card 4/5

Synthesis and Conversions of Sulfur Compounds
of Naphthalene Series Over an Alumina-Silica
Catalyst

782:7
SOV/79-30-3-51/69

The comparative strength of the sulfur bond with
different radicals is shown in scheme 6:



There are 3 tables; and 14 references, 1 U.S., 1 Dutch,
4 German, 8 Soviet. The U.S. reference is: E. D.
Rossini and others, Selected Physical Values and
Thermodynamic Properties of Hydrocarbons and Related
Compounds (1953).

ASSOCIATION: Moscow State University (Moskovskiy gosudarstvennyy
universitet)

SUBMITTED: March 5, 1959
Card 5/5

MINACHEV, Kh.M.; MARKOV, M.A.; LOGINOV, G.A.

Conversions of five- and six-membered cyclanes on rare-
earth oxides. Neftokhimiya 1 no.3:356-361 My-Je '61.
(MIRA 16:11)

1. Institut organicheskoy khimii AN SSSR imeni N.D. Zelinskogo.

5. 1190

31746

S/204/61/001/004/003/005
EO75/E185

AUTHORS: Minachev, Kh.M., Markov, M.A., and Shchukina O.K.

TITLE: Dehydrogenation of cyclohexane on the oxides of rare earth elements

PERIODICAL: Neftekhimiya, v.1, no.4, 1961, 489-493

TEXT: Eight oxides of rare earth elements and yttrium oxide were used as catalysts for the dehydrogenation of cyclohexane. The catalysts were prepared by dissolving the commercial oxides in 27% nitric acid, and precipitating with 12% ammonia solution at 50-60 °C. The dried and washed precipitates were compressed into 4 x 4 mm cylinders and heated at 560 °C in dry air for 8 hours. The surface areas of the oxides so obtained were determined by benzene vapour adsorption. The dehydrogenations were carried out at 515-590 °C under atmospheric pressure. The catalysts were activated at 560 °C for 2 hours by passing through them currents of air, hydrogen or nitrogen. The catalysts with the greatest activity were produced by the activation with nitrogen. Experimental results show that all the catalysts dehydrogenate cyclohexane to benzene.
Card 1/43

Dehydrogenation of cyclohexane ...

³¹⁷⁴⁶
S/204/61/001/004/003/005
E075/E185

The determination of specific areas of the catalysts permitted the calculation of the specific activity and specific coke formation for the various catalysts, and thus their relative overall activities could be compared. The results are given in Table 4. It can be seen that the specific activity and coke formation at 530-560 °C does not differ much inside the yttrium group of the oxides. The yttrium group oxides exceed the cerium oxide group in respect of activity. It was shown that there exists linear dependence between the logarithms of the percentage conversion and the reciprocal temperature of the reaction. The energies of activation calculated from the slopes of the lines had typical values for acidic catalysts in the case of neodymium, gadolinium and holmium oxides, but exceeded 50 kcal for the remaining oxides. There are 4 figures, 4 tables and 8 references: 5 Soviet bloc and 3 non-Soviet-bloc. The English language references read:

Ref. 1: R.A. Briggs, H.S. Taylor,

J. Amer. Chem. Soc., v. 63, 2500, 1941.

Ref. 4: V. I. Komarevsky, Ind. Eng. Chem., v 49, 264, 1957.

Card 2/43

Dehydrogenation of cyclohexane ... ³¹⁷⁴⁶ S/204/61/001/004/003/005
E075/E185

ASSOCIATION: Institut organicheskoy khimii AN SSSR im.
N.D. Zelinskogo
(Institute of Organic Chemistry, AS USSR, imeni
N.D. Zelinskiy)

SUBMITTED: June 7, 1961

Card 3/4

MINACHEV, Kh.M.; MARKOV, M.A.; SHCHUKINA, O.K.

Dehydrocyclization of n-heptane over rare earth oxides. *Neftekhimiya*
1 no.5:610-612 S-O '61. (MIRA 15:2)

1. Institut organicheskoy khimii AN SSSR imeni N.D.Zelinskogo.
(Heptane)(Aromatization)(Rare earth oxides)

3/081/02/000/009/032/075
3158/3101

AUTHORS: Tit's-shvertsova, I. N., Danilova, T. A., Larkov, A. A.,
Stepanova, I. I., Osipenko, Ts. D.

TITLE: Conversion of organosulfur compounds of the α - and β -naphthalene series in the presence of an aluminosilicate catalyst

PERIODICAL: Referativnyy zhurnal. Khimiya, no. 9, 1962, 228, abstract
Zhurnal Obshch. Khimii i Seraorgan. Soedineniy, soderzhashchikh
v neftyakh i nefteproduktakh. v. 4", L., Gostoptekhnizat, 1961,
141 - 144)

TEXT: Contact conversions of organosulfur compounds of naphthalene as carried out at 300°C on an aluminosilicate catalyst under conditions described earlier (Zh. obshch. Khimii, v. 21, 1951, 242) are reexamined. α - and β -thionaphthols (α - and β -I) were synthesized for research, α - and β -naphthyldecylsulfides (α - and β -II) and α - and β -naphthylcyclopentylsulfides (α - and β -III) synthesized for the first time. It was found that under these conditions α -I and β -I are converted to $C_{10}H_8$ and H_2S similarly to the thiophenols studied earlier the respective yields being 52 and 42%.
Card 1/2

Conversion of organosulfur compounds ...

S/081/62/000/039/012/0.5

B158/B101

by weight of catalyst. As established previously (see UCh, zap. ISU, v.11, 1953, 263), in the case of mixed sulfides of the C_6H_5SR type (R being an alkyl or cycloalkyl), the bond between the sulfur and R is always ruptured. In the case of α -II, it was found that $C_{10}H_8$ and $C_{10}H_{21}SH$ are formed with further conversion of the latter to $C_{10}H_{20}$ and H_2S . α -III also decomposes in the same way, forming $C_{10}H_8$ and cyclopentanethiol with subsequent conversion of the latter to dicyclopentylsulfide and H_2S . α -III under these conditions decomposes to α -I, cyclopentane, $C_{10}H_8$ and H_2S . In the case of α -II, α -I, $C_{10}H_{21}SH$, a decene-decane fraction and H_2S were detected. Consequently the bond between the sulfur and the benzene ring in mixed sulfides is much more stable and was not ruptured in any of the cases examined. The bond between the sulfur and the $C_{10}H_8$ in the α -position is far less stable. The bond between the sulfur and the alkyl and naphthyl in the α -position is more stable than that between the sulfur and naphthene rings.

[Abstracter's note: Complete translation.]

Card 2/2

MINACHEV, Kh.M.; MARKOV, M.A.; KHODAKOV, Yu.S.

Effect of gamma rays on the catalytic activity of platinized
aluminosilicate. Izv. AN SSSR. Otd.khim.nauk no.7:1227-1230
Jl '61. (MIRA 14:7)

1. Institut organicheskoy khimii im. N.D. Zelinskogo AN SSSR.
(Aluminosilicates) (Catalysis) (Gamma rays)

S. 000, 000, 000, 000, 000
B. 000, 000, 000, 000, 000

AUTHORS: Minachev, Kh. M., Markov, M. A., and St. Ivanov, D. K.
TITLE: Investigation of the catalytic properties of rare earths
PERIODICAL: Akademiya nauk SSSR. Izvestiya. Otdeleniye khim. nauk, no. 8, 1961, 1507-1511

TEXT: The authors investigated the catalytic properties of rare earths and a mixed catalyst from 15 % erbium oxide and 85 % aluminum oxide. The specific surface of the catalysts was determined by the Brunauer-Emmett-Teller dynamic method by A. M. Rubinshteyn and V. A. Alkhalilov (Bull. Akad. Nauk AN SSSR. Otd. khim. n., 1955, 1296) amounted to 190 m²/g for Er₂O₃/Al₂O₃. The properties of the catalysts are listed in Table 1. The experiments were carried out at temperatures from 545°-590°C and atmospheric pressure with a flow rate of 0.25 hr⁻¹. Before each experiment the catalysts were reduced with hydrogen at 550°C. After the experiment the catalysts were oxidized at 550°C. Card 1/6

Investigation of the catalytic...

500°-520°C with air, which was diluted with nitrogen. The yield of 5-7 %. During the regeneration the amount of catalyst was determined. The experiments showed that the liquid catalysis products were analyzed in a gas liquid chromatograph (Ref. 8: D. A. Kondrat'yev, M. A. Markov and K. M. Markov, *laboratoriya 25*, 1301 (1959)). The analysis was carried out according to the method by G. P. Kaufman (Ref. 9: G. P. Kaufman, *zhurnal khim. fiziki*, 1937). An adsorption chromatograph with thermal conductivity detector was used for the analysis of gaseous products. The experiments showed that the yield of gaseous products during the conversion of cyclohexane on Er_2O_3 increases from 14.5 % at a temperature increase from 545° to 590°C. Simultaneously the yield of gaseous products increases from 14.5 % to 16.5 % and the yield of liquid products from 0.5 % to 1.2 %. During the conversion of cyclohexane on Er_2O_3 the yield of liquid catalysates amounted to 98 % at 545°C and 97 % at 590°C. The yield of gaseous products increased from 14.5 % to 16.5 % at suitable temperatures. The amount of coke accumulated on the catalyst was about equal in both cases. During the conversion of cyclohexane on Er_2O_3

Card 2/6

Investigation of the catalytic....

S/062/51 000/006, 008/010
B117/B204

hydrocarbons, the yield of liquid catalysates on pure Er_2O_3 was much higher than on $\text{Er}_2\text{O}_3/\text{Al}_2\text{O}_3$. However, due to the lower stability of n-heptane and ethyl cyclopentane it was not so big as for cyclohexane. Pure Er_2O_3 thus has a much weaker cracking effect than $\text{Er}_2\text{O}_3/\text{Al}_2\text{O}_3$. The analysis of gaseous catalysis products showed that at any rate the gas obtained on $\text{Er}_2\text{O}_3/\text{Al}_2\text{O}_3$ is richer in hydrocarbons than that obtained on Er_2O_3 . During the conversion of cyclohexane and n-heptane the amount of saturated and unsaturated hydrocarbon is about equal. In the catalysis of ethyl cyclopentane, the gas produced on $\text{Er}_2\text{O}_3/\text{Al}_2\text{O}_3$ contains 14.1 % unsaturated and 10.4 % saturated hydrocarbons. The analysis of the liquid catalysis products showed that the product obtained during the conversion of cyclohexane consists of unchanged hydrocarbons, benzene, methyl cyclopentane and cyclohexane. The benzene content in the product obtained on $\text{Er}_2\text{O}_3/\text{Al}_2\text{O}_3$ amounted to 4.1 % at 545°C and 22.6 % at 590°C. On pure Er_2O_3 , the benzene content at 545°C was 1.3 % and at 590°C 11.1 %. The

Card 3/6

Investigation of the catalytic...

S. 122, 123, 124, 125
B. 122, 123, 124, 125

unchanged n-heptane, toluene and benzene was identified during the conversion of n-heptane. The content of aromatic hydrocarbons was about equal in the liquid catalysate in the presence of both catalysts. The yield of liquid catalysate, however, was much higher on pure Er_2O_3 than on $\text{Er}_2\text{O}_3/\text{Al}_2\text{O}_3$. More aromatic hydrocarbons are thus obtained on Er_2O_3 per hydrocarbon used than on Er_2O_3 . The presence of benzene in the catalysis products of n-heptane points towards the demethylation process. When passing through ethyl cyclopentane, no toluene was established on Er_2O_3 . In the product obtained on $\text{Er}_2\text{O}_3/\text{Al}_2\text{O}_3$, however, 0.1% toluene were found. The iodine numbers of the products obtained during the conversion of cyclohexane and n-heptane on both catalysts were usually not higher than 15. The iodine number of the product obtained from cyclohexane on the mixed catalyst at 590°C , i.e. 29, was an exception. During catalysis of ethyl cyclopentane, the iodine numbers of the catalysates were 29 on Er_2O_3 and 52 on $\text{Er}_2\text{O}_3/\text{Al}_2\text{O}_3$. There are 11 references, 1 table and 10 references: 6 Soviet-bloc and 4 non-Soviet-bloc. The title:

Card 4/6

Investigation of the catalytic...

S/062/61/000/008/008/010.
B117/B206

references to English-language publications read as follows: R. A. Briggs,
H. S. Taylor, J. Amer. Chem. Soc. 63, 2500 (1941); V. I. Komarevsky,
Industr. and Engng. Chem. 49, 264 (1957); g. E. Green, Nature 180,
N 4580, 295 (1957).

ASSOCIATION: Institut organicheskoy khimii im. N. D. Zelinskogo Akademii
nauk SSSR (Institute of Organic Chemistry imeni
N. D. Zelinskiy, AS USSR) ✓

SUBMITTED: December 13, 1960

Card 5/6

5 1190

2209, 1274 1275

27493

S/062/61/000/009/008/014

B117/B101

AUTHORS: Minachev, Kh. M., Markov, M. A., and Shchukina, O. K.

TITLE: Study of the catalytic properties of rare earth oxides
2. Transformation of cyclohexene, 1-methyl cyclohexene-1, and n-heptene-1 on erbium oxide

PERIODICAL: Akademiya nauk SSSR. Izvestiya. Otdeleniye khimicheskikh nauk, no. 9, 1961, 1665-1669

TEXT: The present work which continues the studies on the catalytic properties of rare earth oxides was undertaken with a view to investigating the behavior of several cycloalkenes on pure erbium oxide and a mixed catalyst (15% erbium oxide, 85% aluminum oxide) at 510°-515°C. Catalyst preparation, reaction conditions and method of analyzing the catalyst have been described previously (Ref. 1: Izv. AN SSSR. Otd. khim. n. 1961, no. 8). The following hydrocarbons were used: cyclohexene; 1-methyl cyclohexene-1, n-heptene-1; the yields of liquid catalysis products obtained by passing the two first-mentioned cycloalkenes over Er_2O_3 were independent of time and amounted to 80.7% and 83.4%, respectively. They contained no hydro-

Card 1/6

27493

S/062/61/000/009/008/001
B117/B101

Study of the catalytic properties ...

carbons of molecular weight higher than that of the initial substance. Chromatographic analysis showed that the gas formed simultaneously is made up of 90-96% hydrogen and 4-10% C_1 to C_4 hydrocarbons. Carbonization on Er_2O_3 was insignificant. The product obtained from cyclohexene over Er_2O_3 consisted of benzene (20.23%) and cyclohexene. In experiments using 1-methyl cyclohexene-1 and the same catalyst, 1-methyl cyclohexene-1, a mixture of 1-methyl cyclohexene-2 and 1-methyl cyclohexene-3, and toluene were identified in the reaction product. The results obtained in the conversion of 1-methyl cyclohexene-1 on Er_2O_3 are given in Table 2. The tests with cyclohexene and 1-methyl cyclohexene-1 over Er_2O_3/Al_2O_3 showed that the yield of liquid products increases with time. The gases analyzed consisted of 85%-93% hydrogen and 7-15% C_1 to C_4 hydrocarbons. As in the case of Er_2O_3 , the gas was richer in hydrocarbons towards the beginning of the experiments. Carbonization on Er_2O_3/Al_2O_3 was higher than on pure erbium oxide. The composition of the catalyzate obtained from cyclohexene over Er_2O_3/Al_2O_3 is represented in Table 3. The composition of the

Card 2/6

27493

S/062/61/000/003/008/014

B117/B101

Study of the catalytic properties ...

catalyzate obtained from 1-methyl cyclohexene-1 was only partially clarified, since the chromatogram was greatly complicated by the formation of dimethyl cyclopentenenes. A sample drawn within the first two hours was found to contain 63% toluene. The total yield of catalysis products in tests with n-heptene-1 over Er_2O_3 was 74.5% and over $\text{Er}_2\text{O}_3/\text{Al}_2\text{O}_3$ 68.3%. The gas formed over $\text{Er}_2\text{O}_3/\text{Al}_2\text{O}_3$ contained double the quantity of hydrocarbons that was obtained over Er_2O_3 . The liquid catalysis products contained 6% toluene in the case of Er_2O_3 and 8.5% in the case of $\text{Er}_2\text{O}_3/\text{Al}_2\text{O}_3$. The chromatogram of the product obtained over $\text{Er}_2\text{O}_3/\text{Al}_2\text{O}_3$ exhibited six peaks in the C_7 hydrocarbons region, apart from toluene, as compared to two peaks in the case of Er_2O_3 . Er_2O_3 catalysis yielded products containing 91.8% unsaturated hydrocarbons and $\text{Er}_2\text{O}_3/\text{Al}_2\text{O}_3$ 47.0%. In conclusion, the investigation of these two catalysts yielded the following results: They differ inasmuch as the mixed catalysts produced isomerization of the 6-membered ring to a 5-membered ring, whereas this isomeriza-

Card 3/6

27493

S/062/61/000/009/006/011
B117/B101

Study of the catalytic properties . .

tion does not occur with pure erbium oxide. Both catalysts dehydrogenate the tested cycloalkenes to corresponding aromatic hydrocarbons and cause shifting of the double bond in the ring. There are 3 figures, 3 tables and 5 references: 3 Soviet and 2 non-Soviet. The two references to English-language publications read as follows: F. G. Rossini, K. S. Pitzer, R. L. Arnett, R. M. Braun, G. C. Pimentel, Selected values of physical and thermodynamic properties of hydrocarbons and related compounds, Carnegie Press, 1953; E. Gil-av, J. Herling, J. Shabtai, Chem and chem. Ind. no. 9, 1483 (1957).

ASSOCIATION: Institut organicheskoy khimii im. N. D. Zelinskogo Akademii nauk SSSR (Institute of Organic Chemistry imeni N. D. Zelinskiy of the Academy of Sciences USSR)

SUBMITTED: December 13, 1960

Card 4/6

LUKINA, M.Yu.; ZOTOVA, S.V.; MARKOV, M.A.; OVODOVA, V.A.; KAZANSKIY, P.A.,
akademik

Transformations of isopropenylcyclopropane in the presence of
kieselguhr. Dokl. AN SSSR 139 no.2:381-384 J1 '61. (MIRA 14:7)

1. Institut organicheskoy khimii im. N.D. Zelinskogo AN SSSR.
(Propene) (Kieselguhr)

S/204/62/002/002/001/007
I060/I242

AUTHORS: Minachev, Kh.M., Markov, M.A., and Bogomolov, V.I.

TITLE: Conversion of cyclohexane and n alkanes on rare earth oxides

PERIODICAL: Neftekhimiya, v.2, no.2, 1962, 144-149

TEXT: This work is a continuation of the investigation of the catalytic effect of oxides of rare earth elements on dehydrogenation and dehydrocyclization of hydrocarbons. Neodymium, holmium, and yttrium oxides on activated carbon have shown a high activity in the dehydrogenation of cyclohexane and in dehydrocyclization of n alkanes. When the concentration of Nd_2O_3 in the catalyst increases from 0.5 to 5%, the output of benzene from cyclohexane at 500° grows from 19.6 to 49.4%, while higher concentrations of oxide up to 15% do not produce

Card 1/2

S/204/62/002/002/001/007
I060/I242

Conversion of cyclohexane...

any noticeable increase of output of benzene. The gaseous products of catalysis were analyzed in a XT-2M (KhT-2M) unit; the liquid products were analyzed by gas-liquid chromatography. There are 4 figures and 4 tables. ↓

ASSOCIATION: Institut organicheskoy khimii AN SSSR im. N.D. Zelinskogo (Institute of Organic Chemistry im. N.D. Zelinskiy, AS USSR)

SUBMITTED: January 25, 1961

Card 2/2

ACCESSION NR: AT4035162

S/0000/63/000/000/0125/0130

AUTHOR: Minachev, Kh. M.; Markov, M. A.

TITLE: Investigation of the catalytic properties of the rare-earth elements in the transformation of hydrocarbons

SOURCE: AN SSSR. Institut geokhimii i analiticheskoy khimii. Redkozemel'nyye elementy* (Rare-earth elements). Moscow, Izd-vo AN SSSR, 1963, 125-130

TOPIC TAGS: rare earth, catalytic dehydrogenation, rare earth oxide, cyclohexane, n-heptane, ethylcyclopentane, methyl-cyclopentene, methylcyclohexene, n-heptene, dehydrogenation

ABSTRACT: In a series of experiments on the catalytic activity of the rare earth oxides, oxides of La, Nd, Sm, Gd, Ho, Er, Tu, Yb and Y were used as catalysts in the dehydrogenation of cyclohexane, n-heptane, ethylcyclopentane, 1-methylcyclopentene-1, 1-methylcyclohexene-1 and n-heptene-1. The dehydrogenation of cyclohexane was first investigated at 515-580C. Determination of the specific surface of the investigated oxides made it possible to compare their specific catalytic activity and specific coke-forming power during the reaction. Calculation showed that the activation energy for Nd_2O_3 , Gd_2O_3 and Ho_2O_3 are close to the values of typical oxide catalysts, while for other oxides the values exceed 50 kcal/mole. Similar

Card 1/2

ACCESSION NR: AT4035162

results were obtained with n-heptane and cycloolefins. The isomerization of 1-methylcyclopenten-1 and 1-methylcyclohexen-1 is plotted against temperature, and the production of benzene from the latter is compared with different catalysts. Finally, the reduction of cyclohexane and n-heptane is studied over oxides of the rare-earth elements adsorbed onto activated charcoal. Such catalysts were found to be significantly more effective than the oxides alone. Orig. art. has: 7 figures and 2 tables.

ASSOCIATION: Institut geokhimii i analiticheskoy khimii AN SSSR (Institute of Geochemistry and Analytical Chemistry, AN SSSR)

SUBMITTED: 31Oct63

DATE ACQ: 30Apr64

ENCL: 00

SUB CODE: IC, OC

NO REF SOV: 008

OTHER: 003

Cord

2/2

MINACHEV, Kh.M.; MARKOV, M.A.; BOGOMOLOV, V.I.; ENGLINA, F.E.

Transformation of cyclic alcohols on neodymium oxide. Izv.AN
SSSR. Ser.khim. no.1:13-17 Ja '64. (MIRA 17:4)

1. Institut organicheskoy khimii im. N.D.Zelinskogo AN SSSR.

DERBENTSEV, Yu.I.; MARKOV, M.A.; ISAGULYANTS, G.V.; MINACHEV, Kh.M.;
BALANDIN, A.A., akademik; Prinimala uchastiye SHCHUKINA, O.K.

Mechanism of cyclohexane dehydrogenation over holmium oxide studied
with the use of radiocarbon C¹⁴. Dokl. AN SSSR 155 no.1:128-131
Mr '64. (MIRA 17:4)

1. Institut organicheskoy khimii im. N.D.Zelinskogo AN SSSR.

MINACHEV, Kh.M.; MARKOV, M.A.

Dehydrogenation of cyclohexane on rare-earth oxides deposited
on high-ash coals. Izv. AN SSSR. Ser. khim. no.9:1680-1682 '65.
(MIRA 18:9)

1. Institut organicheskoy khimii im. N.D. Zelinskogo AN SSSR.

197 4th 7th 8th 9th 10th 11th 12th 13th 14th 15th 16th 17th 18th 19th 20th 21st 22nd 23rd 24th 25th 26th 27th 28th 29th 30th 31st 32nd 33rd 34th 35th 36th 37th 38th 39th 40th 41st 42nd 43rd 44th 45th 46th 47th 48th 49th 50th 51st 52nd 53rd 54th 55th 56th 57th 58th 59th 60th 61st 62nd 63rd 64th 65th 66th 67th 68th 69th 70th 71st 72nd 73rd 74th 75th 76th 77th 78th 79th 80th 81st 82nd 83rd 84th 85th 86th 87th 88th 89th 90th 91st 92nd 93rd 94th 95th 96th 97th 98th 99th 100th 101st 102nd 103rd 104th 105th 106th 107th 108th 109th 110th 111th 112th 113th 114th 115th 116th 117th 118th 119th 120th 121st 122nd 123rd 124th 125th 126th 127th 128th 129th 130th 131st 132nd 133rd 134th 135th 136th 137th 138th 139th 140th 141st 142nd 143rd 144th 145th 146th 147th 148th 149th 150th 151st 152nd 153rd 154th 155th 156th 157th 158th 159th 160th 161st 162nd 163rd 164th 165th 166th 167th 168th 169th 170th 171st 172nd 173rd 174th 175th 176th 177th 178th 179th 180th 181st 182nd 183rd 184th 185th 186th 187th 188th 189th 190th 191st 192nd 193rd 194th 195th 196th 197th 198th 199th 200th 201st 202nd 203rd 204th 205th 206th 207th 208th 209th 210th 211th 212th 213th 214th 215th 216th 217th 218th 219th 220th 221st 222nd 223rd 224th 225th 226th 227th 228th 229th 230th 231st 232nd 233rd 234th 235th 236th 237th 238th 239th 240th 241st 242nd 243rd 244th 245th 246th 247th 248th 249th 250th 251st 252nd 253rd 254th 255th 256th 257th 258th 259th 260th 261st 262nd 263rd 264th 265th 266th 267th 268th 269th 270th 271st 272nd 273rd 274th 275th 276th 277th 278th 279th 280th 281st 282nd 283rd 284th 285th 286th 287th 288th 289th 290th 291st 292nd 293rd 294th 295th 296th 297th 298th 299th 300th 301st 302nd 303rd 304th 305th 306th 307th 308th 309th 310th 311th 312th 313th 314th 315th 316th 317th 318th 319th 320th 321st 322nd 323rd 324th 325th 326th 327th 328th 329th 330th 331st 332nd 333rd 334th 335th 336th 337th 338th 339th 340th 341st 342nd 343rd 344th 345th 346th 347th 348th 349th 350th 351st 352nd 353rd 354th 355th 356th 357th 358th 359th 360th 361st 362nd 363rd 364th 365th 366th 367th 368th 369th 370th 371st 372nd 373rd 374th 375th 376th 377th 378th 379th 380th 381st 382nd 383rd 384th 385th 386th 387th 388th 389th 390th 391st 392nd 393rd 394th 395th 396th 397th 398th 399th 400th 401st 402nd 403rd 404th 405th 406th 407th 408th 409th 410th 411th 412th 413th 414th 415th 416th 417th 418th 419th 420th 421st 422nd 423rd 424th 425th 426th 427th 428th 429th 430th 431st 432nd 433rd 434th 435th 436th 437th 438th 439th 440th 441st 442nd 443rd 444th 445th 446th 447th 448th 449th 450th 451st 452nd 453rd 454th 455th 456th 457th 458th 459th 460th 461st 462nd 463rd 464th 465th 466th 467th 468th 469th 470th 471st 472nd 473rd 474th 475th 476th 477th 478th 479th 480th 481st 482nd 483rd 484th 485th 486th 487th 488th 489th 490th 491st 492nd 493rd 494th 495th 496th 497th 498th 499th 500th 501st 502nd 503rd 504th 505th 506th 507th 508th 509th 510th 511th 512th 513th 514th 515th 516th 517th 518th 519th 520th 521st 522nd 523rd 524th 525th 526th 527th 528th 529th 530th 531st 532nd 533rd 534th 535th 536th 537th 538th 539th 540th 541st 542nd 543rd 544th 545th 546th 547th 548th 549th 550th 551st 552nd 553rd 554th 555th 556th 557th 558th 559th 560th 561st 562nd 563rd 564th 565th 566th 567th 568th 569th 570th 571st 572nd 573rd 574th 575th 576th 577th 578th 579th 580th 581st 582nd 583rd 584th 585th 586th 587th 588th 589th 590th 591st 592nd 593rd 594th 595th 596th 597th 598th 599th 600th 601st 602nd 603rd 604th 605th 606th 607th 608th 609th 610th 611th 612th 613th 614th 615th 616th 617th 618th 619th 620th 621st 622nd 623rd 624th 625th 626th 627th 628th 629th 630th 631st 632nd 633rd 634th 635th 636th 637th 638th 639th 640th 641st 642nd 643rd 644th 645th 646th 647th 648th 649th 650th 651st 652nd 653rd 654th 655th 656th 657th 658th 659th 660th 661st 662nd 663rd 664th 665th 666th 667th 668th 669th 670th 671st 672nd 673rd 674th 675th 676th 677th 678th 679th 680th 681st 682nd 683rd 684th 685th 686th 687th 688th 689th 690th 691st 692nd 693rd 694th 695th 696th 697th 698th 699th 700th 701st 702nd 703rd 704th 705th 706th 707th 708th 709th 710th 711th 712th 713th 714th 715th 716th 717th 718th 719th 720th 721st 722nd 723rd 724th 725th 726th 727th 728th 729th 730th 731st 732nd 733rd 734th 735th 736th 737th 738th 739th 740th 741st 742nd 743rd 744th 745th 746th 747th 748th 749th 750th 751st 752nd 753rd 754th 755th 756th 757th 758th 759th 760th 761st 762nd 763rd 764th 765th 766th 767th 768th 769th 770th 771st 772nd 773rd 774th 775th 776th 777th 778th 779th 780th 781st 782nd 783rd 784th 785th 786th 787th 788th 789th 790th 791st 792nd 793rd 794th 795th 796th 797th 798th 799th 800th 801st 802nd 803rd 804th 805th 806th 807th 808th 809th 810th 811th 812th 813th 814th 815th 816th 817th 818th 819th 820th 821st 822nd 823rd 824th 825th 826th 827th 828th 829th 830th 831st 832nd 833rd 834th 835th 836th 837th 838th 839th 840th 841st 842nd 843

1ST AND 2ND EDITIONS										3RD AND 4TH EDITIONS									
PROCEDURES AND PROPERTIES INDEX																			
SA										A53 X									
<p>5007. Permutation Degeneracy in the Vector Model of the Atom. M. Maslov. <i>Comptes Rendus (Doklady) de l'Acad. des Sciences, U.S.S.R.</i> 2. 9. pp. 108-109, 1955. In German.—It is shown that</p> $\text{identities of the type } s_1, s_2 = - (l_1, l_2)^2 - (l_1, l_2) + \frac{1}{2} \text{ (in which } s_1, s_2 \text{ are}$ <p>spin moments and l_1, l_2 are orbital moments) deduced by van Vleck for two equivalent electrons on the basis of the Pauli principle [see Abstract 5006 (1954)], arise from general permutation theory and are valid for any equivalent electrons.]</p> <p style="text-align: right;">C. B. A.</p>																			
ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION										E-27									
FROM SYMBION										FROM GOWARD									
LONDON 72										LONDON 72									
195500 HIF ONLY GET										COLLECTION									
										LONDON 72 ONLY GET									

C

The Dirac vector model for multiplet spectra. M.
Marino. Physik. Z. Sowjetunion 7, 563-564 (1935). - An
expression is derived for the Hamiltonian function of a set
of equiv. electrons of the form: $H = \sum_{\alpha} P_{\alpha}$, where the
 α are polynomials of the scalar products of the orbital
momenta, and the P_{α} are the Slater integrals over the radial
wave functions. Examples are given for the cases of
equiv. p and d electrons. Morris Munkat

ASB SLA METALLURGICAL LITERATURE CLASSIFICATION

SECOND

(10108) 417 ONV JBI

REVISION

10108 417 ONV JBI

1ST AND 2ND ORDERS																										3RD AND 4TH ORDERS																									
COMMON ELEMENTS													PROCESSING AND PROPERTIES INDEX													1ST AND 2ND ORDERS													3RD AND 4TH ORDERS												
MARKOV, M.																										A53																									
2967. Dirac's Theory of the Electron. Part II. M. Markov.																																																			
Phys. Zeits. d. Sowjetunion, 11: 3, pp. 286-288, 1967. In German.																																																			
The equation of second order for a particle with spin which was suggested																																																			
by the author (see Abstract 2966 (1967)) is treated by a method which																																																			
Pauli and Weisskopf used for the discussion of the equation for a particle																																																			
without spin. The resulting equations contain states of positive as well																																																			
as of negative energy and are therefore useless.																																																			
R. P.																																																			
ASD SLA METALLURGICAL LITERATURE CLASSIFICATION																																																			
1900-1960																																																			
1961-1970																																																			
1971-1980																																																			
1981-1990																																																			
1991-2000																																																			

5322. Quantum Potentials and the Electromagnetic Field. M. Markov. *Phys. Zeits. d. Sowjetunion*, 12, 1, pp. 108-110, 1937. In German.—The author points out that, in his opinion, contradictions are involved in quantized electrodynamics such as those of Pauli-Hellmberg.

G. C. McV.

(Quantum postulate and the concept of the electro-
magnetic field. M. Markov. *J. Exptl. Theoret. Phys.*
S. S. R.) 8, 14 (1938). E. H. Rathmann

Quantum electrodynamics. M. Markov. *J. Exp. Theoret. Phys.* (U. S. S. R.) 8, 1246 (1958). M. considers the limiting conditions necessary in order that the error of measurement very near to the point electron should be of the order of the classically assumed field.
J. H. Rathmann

W

Nonelastic dispersion of photons about nuclei with pair production M. Markov *Compt. rend. acad. sci. R. S. S. 20, 1257 (1968) in English; Math.* By use of Williams method M. derives relations for the cross sections for the processes: 1) pair production in the field of the nucleus by the photon and 2) the non-elastic dispersion of a photon about a nucleus accompanied by pair production. The ratio of the second process to the first is independent of the energy and proportional to the fine structure const.

3

Earl A. Gulbransen

ASD 34.0 METALLURGICAL LITERATURE CLASSIFICATION

147380

10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

MARKOV, M. A.

"Role of the Zero-State of Radiation Oscillators in the Higher Approximations of Quantum Electrodynamics," Dokl. AN, 23, No. 9, 1939. Inst. Physics. Acad. Sci., cl 39.
 "Concerning the Four Dimensional-Expanded Electron Relativistic Quantum Zone" Zhur. Phys., 453, No. 6, Vol. II, 1940
 "On the Relativistically Invariant (Cutting-Off Factor) In Electrodynamics," Dokl. AN, 40, No. 1, 1943. P.N. Lebedev Phys. Inst., Acad. Sci. cl943-.
 "Electrodynamics of Dirac-Fock-Podolsky and the Self-energy of the Electron," Dokl. AN, 40, No. 6, 1943. P.N. Lebedev Phys. Inst, Acad. Sci. cl943-.
 "The Many Body Problem in the Classical Relativistic Theory", Zhur. Phys., 42, No. 1. Vol. VII, 1943
 P. N. Lebedev Physical Institute, Academy of Science of the USSR, cl941-.
 "On the Criterion of Relativistic Invariance," Zhur. Eksper. i Teoret. Fiz., 14, No. 9, 1946. Lebedev Physical Inst., Acad. Sci. USSR, -1946-.

sw 260

Is the "self-field" of the particle a physically observable quantity?
M. Markov (Compt rend Acad Sci U.R.S.S., 1943 61, 14-17)
A general theoretical discussion on the basis of existing theories

DATE 11/1/43

Phys. Inst. im. P. N. Lebedev, AN SSSR.

117 AND 120 (ADDED)

PROCESSES AND PROPERTIES INDEX

Proper masses and magnetic moments of elementary particles and the Wentzel Dirac process. M. Markov (P. N. Lebedev Phys. Inst., Academy of Sciences, U. S. S. R.) *Comput. rend. acad. sci. U. R. S. S. R.* 47, No. 4, 177, Nov 1915, *Doklady Akad. Nauk S. S. S. R.* 47, No. 3, 182, 5 (1943), Math. Certain difficulties in the theory of elementary particles with the field can be overcome by the aid of the α -process proposed by Dirac (C.A. 36, 3725). The real proper mass is the same for all elementary particles and equals that of the electron, which is a universal const. and cannot be reduced to a field. All differences in the masses of elementary particles are due to the interaction with the various fields, which leads to the appearance of new terms in this const. The problem is considered quantitatively. H. G. McCann

COMMON ELEMENTS

OPEN

MATERIALS INDEX

ASB-55A METALLURGICAL LITERATURE CLASSIFICATION

SECTION DIVISION

SECTION HIR ONE DEL

SECTION ONE

SECTION TWO

SECTION THREE

SECTION FOUR

SECTION FIVE

SECTION SIX

SECTION SEVEN

SECTION EIGHT

SECTION NINE

SECTION TEN

SECTION ELEVEN

SECTION TWELVE

SECTION THIRTEEN

SECTION FOURTEEN

SECTION FIFTEEN

SECTION SIXTEEN

SECTION SEVENTEEN

SECTION EIGHTEEN

SECTION NINETEEN

SECTION TWENTY

SECTION TWENTY ONE

SECTION TWENTY TWO

SECTION TWENTY THREE

SECTION TWENTY FOUR

SECTION TWENTY FIVE

SECTION TWENTY SIX

SECTION TWENTY SEVEN

SECTION TWENTY EIGHT

SECTION TWENTY NINE

SECTION THIRTY

SECTION THIRTY ONE

SECTION THIRTY TWO

SECTION THIRTY THREE

SECTION THIRTY FOUR

SECTION THIRTY FIVE

SECTION THIRTY SIX

SECTION THIRTY SEVEN

SECTION THIRTY EIGHT

SECTION THIRTY NINE

SECTION FORTY

SECTION FORTY ONE

SECTION FORTY TWO

SECTION FORTY THREE

SECTION FORTY FOUR

SECTION FORTY FIVE

SECTION FORTY SIX

SECTION FORTY SEVEN

SECTION FORTY EIGHT

SECTION FORTY NINE

SECTION FIFTY

SECTION FIFTY ONE

SECTION FIFTY TWO

SECTION FIFTY THREE

SECTION FIFTY FOUR

SECTION FIFTY FIVE

SECTION FIFTY SIX

SECTION FIFTY SEVEN

SECTION FIFTY EIGHT

SECTION FIFTY NINE

SECTION SIXTY

SECTION SIXTY ONE

SECTION SIXTY TWO

SECTION SIXTY THREE

SECTION SIXTY FOUR

SECTION SIXTY FIVE

SECTION SIXTY SIX

SECTION SIXTY SEVEN

SECTION SIXTY EIGHT

SECTION SIXTY NINE

SECTION SEVENTY

SECTION SEVENTY ONE

SECTION SEVENTY TWO

SECTION SEVENTY THREE

SECTION SEVENTY FOUR

SECTION SEVENTY FIVE

SECTION SEVENTY SIX

SECTION SEVENTY SEVEN

SECTION SEVENTY EIGHT

SECTION SEVENTY NINE

SECTION EIGHTY

SECTION EIGHTY ONE

SECTION EIGHTY TWO

SECTION EIGHTY THREE

SECTION EIGHTY FOUR

SECTION EIGHTY FIVE

SECTION EIGHTY SIX

SECTION EIGHTY SEVEN

SECTION EIGHTY EIGHT

SECTION EIGHTY NINE

SECTION NINETY

SECTION NINETY ONE

SECTION NINETY TWO

SECTION NINETY THREE

SECTION NINETY FOUR

SECTION NINETY FIVE

SECTION NINETY SIX

SECTION NINETY SEVEN

SECTION NINETY EIGHT

SECTION NINETY NINE

SECTION ONE HUNDRED

SECTION ONE HUNDRED ONE

SECTION ONE HUNDRED TWO

SECTION ONE HUNDRED THREE

SECTION ONE HUNDRED FOUR

SECTION ONE HUNDRED FIVE

SECTION ONE HUNDRED SIX

SECTION ONE HUNDRED SEVEN

SECTION ONE HUNDRED EIGHT

SECTION ONE HUNDRED NINE

SECTION TWO HUNDRED

SECTION TWO HUNDRED ONE

SECTION TWO HUNDRED TWO

SECTION TWO HUNDRED THREE

SECTION TWO HUNDRED FOUR

SECTION TWO HUNDRED FIVE

SECTION TWO HUNDRED SIX

SECTION TWO HUNDRED SEVEN

SECTION TWO HUNDRED EIGHT

SECTION TWO HUNDRED NINE

SECTION THREE HUNDRED

SECTION THREE HUNDRED ONE

SECTION THREE HUNDRED TWO

SECTION THREE HUNDRED THREE

SECTION THREE HUNDRED FOUR

SECTION THREE HUNDRED FIVE

SECTION THREE HUNDRED SIX

SECTION THREE HUNDRED SEVEN

SECTION THREE HUNDRED EIGHT

SECTION THREE HUNDRED NINE

SECTION FOUR HUNDRED

SECTION FOUR HUNDRED ONE

SECTION FOUR HUNDRED TWO

SECTION FOUR HUNDRED THREE

SECTION FOUR HUNDRED FOUR

SECTION FOUR HUNDRED FIVE

SECTION FOUR HUNDRED SIX

SECTION FOUR HUNDRED SEVEN

SECTION FOUR HUNDRED EIGHT

SECTION FOUR HUNDRED NINE

SECTION FIVE HUNDRED

SECTION FIVE HUNDRED ONE

SECTION FIVE HUNDRED TWO

SECTION FIVE HUNDRED THREE

SECTION FIVE HUNDRED FOUR

SECTION FIVE HUNDRED FIVE

SECTION FIVE HUNDRED SIX

SECTION FIVE HUNDRED SEVEN

SECTION FIVE HUNDRED EIGHT

SECTION FIVE HUNDRED NINE

SECTION SIX HUNDRED

SECTION SIX HUNDRED ONE

SECTION SIX HUNDRED TWO

SECTION SIX HUNDRED THREE

SECTION SIX HUNDRED FOUR

SECTION SIX HUNDRED FIVE

SECTION SIX HUNDRED SIX

SECTION SIX HUNDRED SEVEN

SECTION SIX HUNDRED EIGHT

SECTION SIX HUNDRED NINE

SECTION SEVEN HUNDRED

SECTION SEVEN HUNDRED ONE

SECTION SEVEN HUNDRED TWO

SECTION SEVEN HUNDRED THREE

SECTION SEVEN HUNDRED FOUR

SECTION SEVEN HUNDRED FIVE

SECTION SEVEN HUNDRED SIX

SECTION SEVEN HUNDRED SEVEN

SECTION SEVEN HUNDRED EIGHT

SECTION SEVEN HUNDRED NINE

SECTION EIGHT HUNDRED

SECTION EIGHT HUNDRED ONE

SECTION EIGHT HUNDRED TWO

SECTION EIGHT HUNDRED THREE

SECTION EIGHT HUNDRED FOUR

SECTION EIGHT HUNDRED FIVE

SECTION EIGHT HUNDRED SIX

SECTION EIGHT HUNDRED SEVEN

SECTION EIGHT HUNDRED EIGHT

SECTION EIGHT HUNDRED NINE

SECTION NINE HUNDRED

SECTION NINE HUNDRED ONE

SECTION NINE HUNDRED TWO

SECTION NINE HUNDRED THREE

SECTION NINE HUNDRED FOUR

SECTION NINE HUNDRED FIVE

SECTION NINE HUNDRED SIX

SECTION NINE HUNDRED SEVEN

SECTION NINE HUNDRED EIGHT

SECTION NINE HUNDRED NINE

SECTION TEN HUNDRED

SECTION TEN HUNDRED ONE

SECTION TEN HUNDRED TWO

SECTION TEN HUNDRED THREE

SECTION TEN HUNDRED FOUR

SECTION TEN HUNDRED FIVE

SECTION TEN HUNDRED SIX

SECTION TEN HUNDRED SEVEN

SECTION TEN HUNDRED EIGHT

SECTION TEN HUNDRED NINE

SECTION ELEVEN HUNDRED

SECTION ELEVEN HUNDRED ONE

SECTION ELEVEN HUNDRED TWO

SECTION ELEVEN HUNDRED THREE

SECTION ELEVEN HUNDRED FOUR

SECTION ELEVEN HUNDRED FIVE

SECTION ELEVEN HUNDRED SIX

SECTION ELEVEN HUNDRED SEVEN

SECTION ELEVEN HUNDRED EIGHT

SECTION ELEVEN HUNDRED NINE

SECTION TWELVE HUNDRED

SECTION TWELVE HUNDRED ONE

SECTION TWELVE HUNDRED TWO

SECTION TWELVE HUNDRED THREE

SECTION TWELVE HUNDRED FOUR

SECTION TWELVE HUNDRED FIVE

SECTION TWELVE HUNDRED SIX

SECTION TWELVE HUNDRED SEVEN

SECTION TWELVE HUNDRED EIGHT

SECTION TWELVE HUNDRED NINE

SECTION THIRTEEN HUNDRED

SECTION THIRTEEN HUNDRED ONE

SECTION THIRTEEN HUNDRED TWO

SECTION THIRTEEN HUNDRED THREE

SECTION THIRTEEN HUNDRED FOUR

SECTION THIRTEEN HUNDRED FIVE

SECTION THIRTEEN HUNDRED SIX

SECTION THIRTEEN HUNDRED SEVEN

SECTION THIRTEEN HUNDRED EIGHT

SECTION THIRTEEN HUNDRED NINE

SECTION FOURTEEN HUNDRED

SECTION FOURTEEN HUNDRED ONE

SECTION FOURTEEN HUNDRED TWO

SECTION FOURTEEN HUNDRED THREE

SECTION FOURTEEN HUNDRED FOUR

SECTION FOURTEEN HUNDRED FIVE

SECTION FOURTEEN HUNDRED SIX

SECTION FOURTEEN HUNDRED SEVEN

SECTION FOURTEEN HUNDRED EIGHT

SECTION FOURTEEN HUNDRED NINE

SECTION FIFTEEN HUNDRED

SECTION FIFTEEN HUNDRED ONE

SECTION FIFTEEN HUNDRED TWO

SECTION FIFTEEN HUNDRED THREE

SECTION FIFTEEN HUNDRED FOUR

SECTION FIFTEEN HUNDRED FIVE

SECTION FIFTEEN HUNDRED SIX

SECTION FIFTEEN HUNDRED SEVEN

SECTION FIFTEEN HUNDRED EIGHT

SECTION FIFTEEN HUNDRED NINE

SECTION SIXTEEN HUNDRED

SECTION SIXTEEN HUNDRED ONE

SECTION SIXTEEN HUNDRED TWO

SECTION SIXTEEN HUNDRED THREE

SECTION SIXTEEN HUNDRED FOUR

SECTION SIXTEEN HUNDRED FIVE

SECTION SIXTEEN HUNDRED SIX

SECTION SIXTEEN HUNDRED SEVEN

SECTION SIXTEEN HUNDRED EIGHT

SECTION SIXTEEN HUNDRED NINE

SECTION SEVENTEEN HUNDRED

SECTION SEVENTEEN HUNDRED ONE

SECTION SEVENTEEN HUNDRED TWO

SECTION SEVENTEEN HUNDRED THREE

SECTION SEVENTEEN HUNDRED FOUR

SECTION SEVENTEEN HUNDRED FIVE

SECTION SEVENTEEN HUNDRED SIX

SECTION SEVENTEEN HUNDRED SEVEN

SECTION SEVENTEEN HUNDRED EIGHT

SECTION SEVENTEEN HUNDRED NINE

SECTION EIGHTEEN HUNDRED

SECTION EIGHTEEN HUNDRED ONE

SECTION EIGHTEEN HUNDRED TWO

SECTION EIGHTEEN HUNDRED THREE

SECTION EIGHTEEN HUNDRED FOUR

SECTION EIGHTEEN HUNDRED FIVE

SECTION EIGHTEEN HUNDRED SIX

SECTION EIGHTEEN HUNDRED SEVEN

SECTION EIGHTEEN HUNDRED EIGHT

SECTION EIGHTEEN HUNDRED NINE

SECTION NINETEEN HUNDRED

SECTION NINETEEN HUNDRED ONE

SECTION NINETEEN HUNDRED TWO

SECTION NINETEEN HUNDRED THREE

SECTION NINETEEN HUNDRED FOUR

SECTION NINETEEN HUNDRED FIVE

SECTION NINETEEN HUNDRED SIX

SECTION NINETEEN HUNDRED SEVEN

SECTION NINETEEN HUNDRED EIGHT

SECTION NINETEEN HUNDRED NINE

SECTION TWENTY HUNDRED

SECTION TWENTY HUNDRED ONE

SECTION TWENTY HUNDRED TWO

SECTION TWENTY HUNDRED THREE

SECTION TWENTY HUNDRED FOUR

SECTION TWENTY HUNDRED FIVE

SECTION TWENTY HUNDRED SIX

SECTION TWENTY HUNDRED SEVEN

SECTION TWENTY HUNDRED EIGHT

SECTION TWENTY HUNDRED NINE

SECTION TWENTY ONE HUNDRED

SECTION TWENTY ONE HUNDRED ONE

SECTION TWENTY ONE HUNDRED TWO

SECTION TWENTY ONE HUNDRED THREE

SECTION TWENTY ONE HUNDRED FOUR

SECTION TWENTY ONE HUNDRED FIVE

SECTION TWENTY ONE HUNDRED SIX

SECTION TWENTY ONE HUNDRED SEVEN

SECTION TWENTY ONE HUNDRED EIGHT

SECTION TWENTY ONE HUNDRED NINE

SECTION TWENTY TWO HUNDRED

SECTION TWENTY TWO HUNDRED ONE

SECTION TWENTY TWO HUNDRED TWO

SECTION TWENTY TWO HUNDRED THREE

SECTION TWENTY TWO HUNDRED FOUR

SECTION TWENTY TWO HUNDRED FIVE

SECTION TWENTY TWO HUND

PA 54791

MARKOV, M.

USSR/Physics
Relativity
Quantum Mechanics

Jul/Aug 1946

"A Certain Criterion of Relativistic Invariance,"
M. Markov, Lebedev Phys Inst, Acad Sci USSR, 8 pp

^{Zhurn. Fiz.}
"Journal of Physics USSR" Vol I, No 4

Attempt to remove field theory difficulties by use
of Hamilton-Jacobi equation, quantum equations of
motion, Dirac's many-time formalism and Heitler-Peng
"calculation scheme," considered from viewpoint of
criterion of relativistic invariance. Received,
19 Aug 1945.

54791

Markov, M.

Markov, M. On the back action of the electromagnetic field on a moving electron. Acad. Sci. USSR, 1946, 10, 159-166 (1946).

The author discusses some one-dimensional solutions of the classical system of equations describing the motion of an extended electron subject to the influence of its own and an external electromagnetic field. The basic equations used are those due to Lorentz. In these equations the electromagnetic fields due to the charge and current density are eliminated by using the results of Belousov [Akad. Nauk SSSR, Zhurnal Eksper. Teoret. Fiz. 9, 658-669 (1939)], who has given a general solution of the Maxwell field equations for arbitrary charges and current densities in terms of Fourier integrals. When this is done one is left with a system of integral equations for the acceleration vector of the center of gravity of the electron. These equations are discussed in the one-dimensional nonrelativistic case, where they reduce to

$$m\ddot{x} = -\frac{4}{3} \frac{e^2}{c} \int_0^\infty \int_0^\infty \dot{x}(s) k \sin kc(t-s) f^2(k) ds dk + F,$$

where m is the inertial mass of the electron, e its charge, c the velocity of light, \dot{x} the acceleration of the electron and F the external force. The function $f(k)$ is determined in terms of the charge density by $f(k) = \int \rho(r) e^{ikr} dr$. The Dirac modification of the classical Lorentz theory of the electron [Proc. Roy. Soc. London, Ser. A, 167, 148-169 (1938); 180, 1-40 (1942); Communications Dublin Inst. Advanced Studies, Ser. A, no. 1 (1943); these Rev. 5, 277; 7, 105] is included in this treatment by setting $f(k) = \cos k\lambda$.

The author then discusses various methods for handling the improper integrals involved and claims that solutions of the form $Be^{i\omega t}$ cannot be obtained if the process of taking the limit $\lambda \rightarrow 0$ is employed at the proper time, that is, after the integral equation is solved. The argument leading to this assertion is not convincing.

A. H. Taub.

Source: Mathematical Reviews,

Vol. 8, No. 3

874 3/29

MARKOV, M.A.

(Markov, M. On a certain violation of relativistic invariance. Acad. Sci. USSR, J. Phys. 10, 333-340 (1946).

If two points in space-time are such that the vector joining them is space-like then the measurement of one physical quantity M at one of these points cannot react on the measurement of a quantity N at the second of these points. That is, the operators representing these quantities at the respective points must commute. The author applies this principle of "causal independence" to the operators representing the vector potential of an electromagnetic field and the Dirac-Hamiltonian for a system of n electrons interacting with this field. He uses the Dirac-Fock-Podolsky many time formalism to obtain a condition on the factors introduced by any "cut-off" process for the Fourier expansion of the vector potential. Introducing such cut-off factors is equivalent to modifying the commutation equations for the Fourier components of the vector potential. When these are written as $[A_{\mu}^+(k), A_{\nu}^-(k')] = c(k, \lambda) \delta_{\mu\nu}$, instead of $[A_{\mu}^+(k), A_{\nu}^-(k')] = \delta_{\mu\nu}$, the function $c(k, \lambda)$ cannot be arbitrary but must satisfy the condition referred to above.

The author shows that, if $c = c^{\mu\nu}$, $\mu = 1, 2, 3, 4$, this condition is violated and hence the principle of causal independence at two relatively space-like points does not hold. He interprets the result as follows. The introduction of the function $c^{\mu\nu}$ in the commutation relations is equivalent to postulating a certain charge distribution for the electron, as may be seen from an application of the many time formalism. Signals are propagated across this charge distribution at a velocity greater than that of light and as a result the causal independence principle is violated. He then states that any choice of an integrable function for $c(k, \lambda)$ would determine a charge distribution and the same difficulty would then arise.

The author then reviews the singular function $c(k, \lambda)$ introduced in the Wentzel-Dirac limiting process and a generalization of this function. He also reviews the Heitler-Peng "calculation scheme" [Proc. Cambridge Philos. Soc. 38, 296-312 (1942)] in light of the causal independence principle and finds that this principle is satisfied.

A. H. Taub (Seattle, Wash.)

Source: Mathematical Reviews,

Vol. 8, No. 5

Markov, M. A. On the criterion of relativistic invariance.

Akad. Nauk SSSR. Zhurnal Eksp. Teor. Fiz. 16, 790-799 (1946). (Russian. English summary.)

[Reviewer's note: the Russian title should be translated

On a criterion of relativistic invariance." Any relativistic

theory based on the notion of space-time continuum ought

to satisfy the following condition. (C) If (t_1, t_2) and (t_1, t_2)

are two world points which cannot be connected by a light

signal, i.e., if the inequality $c|t_1 - t_2| < |x_1 - x_2|$ holds, any

two events at these world points must not influence one

another. The formal relativistic invariance of a theory does

not necessarily imply the condition (C). Using Dirac's

"many-time" formalism the author investigates, from this

point of view, several possible forms of quantum electro-

dynamics (partly generalizing those which have appeared

in the literature). In order to overcome the well-known

convergence difficulties, in most of these theories a four

vector λ is introduced, which is then made to tend to zero.

The author shows that as long as $\lambda \neq 0$ these theories do not

satisfy the condition (C). [Reviewer's note: Since it is

impossible to select a nonvanishing four vector in an invari-

ant fashion, these theories cannot be considered relativis-

tically invariant for $\lambda \neq 0$ unless the components of λ are

introduced as variables which describe new degrees of free-

dom of the system in question. So far only the limiting case

$\lambda \rightarrow 0$ has been assumed to be physically significant.] In the

last section the author briefly discusses the "computational

scheme" of Heitler and Peng [Proc. Cambridge Philos. Soc.

38, 296-312 (1942)] and Heisenberg's S-matrix [Z. Physik

120, 513-538 (1943); these Rev. 4, 292]. A fuller discussion

is reserved for a later publication. [A translation appeared

in Acad. Sci. USSR. J. Phys. 10, 333-340 (1946); these Rev.

8, 303.]

V. Bargmann (Princeton, N. J.).

2

Sum

Source: Mathematical Reviews,

Vol 8 No. 7

Markov, M.

8

Markov, M. On the back action of the electromagnetic
field on a moving electron. Akad. Nauk SSSR Zhurnal
Eksp. Teoret. Fiz. 16, 800-810 (1946). (Russian
English summary)
An English translation is reviewed above.

Source: Mathematical Reviews,

Vol. 8, No. 3

Handwritten signature

C 2
1951

Higher approximations
3A

Higher approximations in the perturbation theory of scalar meson field interaction. M. Mashkov. *Dokl. Akad. Nauk SSSR*, Moscow, 1951, No. 1, 1951. While the 2nd approximation gives Coulomb's law and the Yukawa potential, higher approximations give terms of opposed signs the sum of which becomes zero on each following approximation. The quantum exchange holds also in the 2nd approximation which thus is not the first non vanishing approximation. N. Thon.

MARKOV, M.

Letter to the editor on the limiting λ -process. *Zhur. eksp. i teor. fiz.* 17
no.9:848 '47. (MLRA 6:7)

1. Fizicheskiy institut im. P.N. Lebedeva Akademii Nauk SSSR.
(Quantum theory)

MARKOV, M. A.

2000

Markov, M. A. A classical analogue of the quantum theory of perturbation. *Akad. Nauk SSSR, Zhurnal Eksperimental'noi Fiziki* 18: 510-514 (1918). (Russian)
The Hamilton-Jacobi equation of a classical relativistic system containing charged particles is solved in an iterative approximation by assuming Hamilton's principal function to be expressible as a series in ascending powers of the charge. This procedure is analogous to the perturbation method generally used in the quantum theory of radiation. The method is applied to the problem of the transverse or magnetic self-energy of an electron and the interaction of two fast electrons. The result obtained for the self-energy corresponds to that obtained by Wale, from the older form of Dirac theory. In this case quantum methods in the positron theory have already provided a more desirable (less divergent) result. The interaction formula is not completely worked out, but is stated to correspond to that of Moller, as would be expected.
W. H. Furry

Source: Mathematical Reviews.

Vol. 14, No. 1

1009 Self-Energy of a Moving Charge. M. A. Markov.
Zhur. Ekspl. i Teoret. Fiz. 18, 1130-37 (1968) (in
Russian).

The expressions that are usually given for the self-energy
of a moving charge exhibit an explicit dependence on the
velocity that stands in contradiction to the transformational
properties of the magnitudes involved. It is shown that this
fact is closely related to the presence in the said expres-
sions of diverging integrals. By simple operations the usual
expressions can be given a form that reflects the correct
dependence on the velocity. (auth)

MARKOV, M. A.

PA 19/49T109

USSR/Nuclear Physics - Radioactivity Oct 48
Nuclear Physics - Neutrino

"Theory of the Beta Decomposition in the Case of a Low Upper Boundary for the Beta-Spectrum and the Neutrino Mass," M. A. Markov, Phys Inst imeni P. N. Lebedev, Acad Sci USSR, 4 pp

"Zhur Eksper i Teoret Fiz" Vol XVIII, No 10

Examines beta decomposition assuming that neutrino mass is not zero. In cases of low upper beta-spectrum boundary, mass of neutrino plays essentially different role in variants of beta-decomposition theory. Results are illustrated using H^3 as example. Submitted 20 Apr 48.
19/49T109

MARKOV, M.

USSR/Nuclear Physics - Electrons, Pair
Theory
Nuclear Physics - Neutrettos

Feb 49

"The Pair Theory of Nuclear Forces," W. Lomsadze, M. Markov, Moscow State U,
Phys Inst imeni P. N. Lebedev, Acad Sci USSR, 3 pp

"Zhur Eksper i Teoret Fiz" Vol XIX, No 2

Shows that the neutretto and antineutretto lead to different laws of reaction in the
theory of pair nuclear forces (spin of neutretto $\frac{1}{2}$)

PA32/49T76

USSR/Nuclear Physics - Mesotrons, Apr 49
Capture by Nuclei
Nuclear Physics - Mesotrons, Pair
Production

"The Capture of Negative Mesons by Atomic Nuclei
From the Standpoint of the Pair Theory of Nuclear
Forces," V. Lebedev, M. Markov, Moscow State
U; Phys Inst imeni P. N. Lebedev, Acad Sci USSR,
5 pp

"Zhur, Eksp. i Teoret Fiz." Vol XII, No 4

Attempts to clarify experimental data on meson
capture by atomic nuclei using the pair

38/497101

USSR/Nuclear Physics - Mesotrons, Apr 49
Capture by
Nuclei (Contd.)

theory of meson forces. Concludes that negative
meson will be captured by Fe56, Al27, and S32
nuclei with probability greater than the prob-
ability of spontaneous disintegration of the meson
in the energy interval from -2 Mev to 0. The
capture of mesons by Cl2 and Be7 nuclei for the
same stipulations is absolutely forbidden. Sub-
mitted 10 Jul 48.

38/497101

PA 38/497101

1ST AND 2ND COLUMNS										3RD AND 4TH COLUMNS									
PROCESSES AND PROPERTIES INDEX																			
<p><i>N</i></p> <p><i>8</i></p> <p>MESON EFFECTS ON DEUTERIUM, trans. from Doklady Akad. Nauk S.S.S.R. 75, No. 5, 855-8(1950) M. Markov. 1950. 6p. (NP-2378)</p> <p>The creation of neutral, scalar, and pseudoscalar mesons by capture of fast neutrons by protons, that is, deuterium formation, has been considered for various widths of the potential well and for the case when the potential well equals zero. Plots have been made showing the dependence of cross section on neutron energy.</p>																			
<p>ASB 514 METALLURGICAL LITERATURE CLASSIFICATION</p>																			

MARKOV, M. A., DRABNIKA, S. I., SUVOROV, S. G. (Editor), AKHLAMOV, S. N. (Tech. Editor),
SOKOLOV, A. A. and FEYNBERG, Ye. A.

D. I. Blokhintsev, "Fundamentals of Quantum Mechanics", Osnovy Kvantovoy
Mekhaniki, State Press for Technical-Theoretical Literature.

Table of Contents W-17671, 5 Apr 1951

Markov, M. A. On nonlocalized fields. *I. Akad. Nauk SSSR. Zhurnal Eksper. Teoret. Fiz.* 21, 11-15 (1951). (Russian)

A form of nonlocalizable theory is proposed, in which the component $A_\mu(k)$ of the electromagnetic potential operator with propagation-vector k does not commute with the space-time coordinate x_μ . The commutation law $[A_\mu(k), x_\nu] = -i\delta_{\mu\nu} A_\mu(k)$ is suggested as a possibility, with b a universal constant with the dimension of a length. This theory has some similarity to the nonlocal formalism of H. Yukawa [Physical Rev. (2) 76, 300-301 (1949)]. The consequences of the theory are not explored in detail. The author claims that his formalism gives an unambiguous and consistent definition of a wave-function describing the behaviour of interacting particles so long as these particles are well separated from each other, and in particular including an account of the asymptotic behaviour of the particles at infinity. He states also that the wave-function automatically vanishes when the particles are close together (in comparison with the natural unit of length b) thus forbidding a strictly local description of collision processes.

F. J. Dyson (Ithaca, N. Y.)

Source: Mathematical Reviews,

Vol. 12 No. 10

MARKOV, M. A.

Markov, M. A. On the differences between particles and antiparticles obeying Dirac's equation. Akad. Nauk SSSR. Zhurnal Eksp. Teoret. Fiz. 21: 761-769 (1951). (Russian)

The discussion is organized under the headings: 1) Neutral fields: (a) meson fields, (b) neutrino fields, (c) differences between neutrino and antineutrino fields, (d) charged fields: (a) meson fields, (b) β -forces. The Dirac equation for a nucleon in interaction with one of the above fields is contrasted with that for the antinucleon. Particular attention is devoted to the sign of the specific charge for the five standard transformation types of interaction in order to pick out the "Majorana theories", that is, theories "without antiparticles". The following is typical of the type of result obtained by rather obvious considerations: The specific charge for neutral scalar or pseudovector interaction with antinucleons has the same sign as for that with nucleons, whereas it has the opposite sign in the case of pseudoscalar, vector or tensor interaction. A. J. Coleman (Toronto, Ont.).

Source: Mathematical Reviews,

Vol 13 No. 4

[Handwritten signature]

MARKOV, M. A.

USSR:

123. A dynamically determinable form-factor of elementary particles. M. A. MARKOV. *Zh. eksp. teor. fiz.* 25, No. 3(11) 327-38 (1953) in Russian.

Usually form-factors are assumed to be independent of the forces acting on the particle, which is equivalent to the assumption of a perfectly rigid structure (apart from the kinematic Lorentz contraction). The signal velocity inside such a structure is infinite, in contradiction with the principle of relativity. The usual form-factor theories must, therefore, contain hidden inconsistencies which are responsible for their failure; the nature of these is illustrated in the Tomonaga-Schwinger as well as the S-matrix formulations of rigid form-factor theories. The description of a system with a deformable form-factor $F(x)$ leads to the treatment of $F(x)$ as a new field. It is natural to attempt to identify this field with currently studied fields. From this point of view the Dirac equation for a "free" proton should be regarded as an approximate equation for the motion of the centre

of gravity of a distribution of charge, whose form-factor $F(x)$ is the density $\rho(x)$ of a π -meson field ($\rho(x) = \frac{1}{2}(\dot{\phi}^2 - \phi^2)$ for scalar mesons). The pion field is described by a Klein-Gordon equation which, in turn, contains a form factor $f(x)$ (e.g. $f = \psi^2$, where ψ is the nucleon field). The resulting complicated system of inter-related equations can be approached in a phenomenological way by the use of a simple deformable form-factor. An example is given by taking for F an expression of the type: \exp (minus the scalar product of the nucleon and meson wave-numbers). The resulting modification of the differential cross-section $\sigma(E, \theta)$ for the scattering of π^+ mesons by a proton is given by a simple function of E and θ , which predicts an eventual decrease of σ with E and, for θ' (but not θ), a relatively augmented scattering at large angles. Similar considerations lead to simple predictions regarding the modifications in the photo-production of charged and neutral mesons. W. Z. SWIATECKI

MARKOV, M. A.

USSR/Nuclear Physics - Particle Theory

Nov 53

"Nonlocal Fields and the Complex Nature of the 'Elementary' Particles (the Dynamically Deformable Form-Factor)," M. A. Markov

Usp Fiz Nauk, Vol 51, No 3, pp 317-341

Reviews existing attempts, Western and Soviet, to consider the elementary particles as fields (extensions). Discusses the phenomena appearing in the case of the approximate applicability of nondeformable form-factors (the creation of charged mesons by photons, and the

272T53

photo-generation of neutral mesons). Cites 8 Western and 3 Soviet references (D. Blokhintsev, ZhETF 18, 1948; M. Karkov, ZhETF, 10, 1940, and 21, 1951).

SECRET

1. The following information was obtained from a source who has provided reliable information in the past and is being provided to you for your information.

MARKOV, M. A.

"On the Classification of Fundamental Particles," published by the Academy of Sciences
USSR, 1955

English translation available Branch #5 E-3672

Paper presented at Rochester, N.Y. Conference on High Energy Nuclear Physics, 1955

FD-2354

USSR/Nuclear Physics - Hyperions and mesons

Card 1/2 Pub. 146 - 19/34

Author : Markov, M., and Stakhanov, V.

Title : Possible beta-decay of hyperions and k-mesons

Periodical : Zhur. eksp. i teor. fiz. 28, (40, Jun 1955

Abstract : The writers of the present note consider as an obvious established fact that Λ^0 -particles can enter the composition of complex nuclei on par with nucleons. Unknown, however, is how far the kinship of hyperions and nucleons extends (V. Stakhanov, Diploma work, Moscow State University, 1954), in every case there being certain grounds for considering the Λ^0 -particle as a nucleon situated in a certain excited state with all the consequences from this notion (V. Stakhanov, ibidem); e. g. such a nucleon could be beta-active: $\Lambda^0 \rightarrow p + e^- + \bar{\nu}$. The purpose of the present note is to turn attention to the fact that in this case thanks to the high upper bound of the energy of decay one can expect small lifetimes (tau sub-beta) comparable with the observed lifetimes of Λ^0 -particles, i. e. sub-pi ($\Lambda^0 \rightarrow \pi^- + p$). The authors note that these ideas can be

Card 2/2

FD-2354

applied also to the case of K-mesons (M. A. Markov, Dok. Akad. Nauk SSR, 101, 449, 1955) under certain conditions (zero spin, decay with emission of Fermi particles, etc.). Three references: e. g. W. Fry and M. Swami, Phys. Rev. 96, 1954.

Institution : Physical Institute im. Lebedev, Acad. Sci. USSR

Submitted : Feb 9, 1955

MARKOV, M. A.

✓ The theory of Λ^0 particles. P. S. Ince and M. A. Markov
P. N. Lebedev Phys. Inst., Acad. Sci. U.S.S.R., Moscow)
Zhur. Eksp. i Teor. Fiz. 29, 111-14 (1955). There is
examined methodically the possibility of one class of equa-
tions with internal degrees of freedom for the description of
the Λ^0 particles as excited state of the nucleon.
Werner Jacobson

①

Markov, M.



(Markov, M. On the theory of the dynamically deformable form-factor. Dokl. Akad. Nauk SSSR (N.S.) 161, 1551-54 (1955). (Russian)

This is a very general and preliminary discussion of methods of making interactions between elementary particles non-local, without introducing rigid structures defined by fixed form-factors. The proposal is to make the "form-factor" itself a dynamical part of the system, reacting to external forces in a causal way. A set of equations partially embodying this idea is presented.

F. J. Dyson (Princeton, N.J.).

1 - P/W

MARKOV, M.A.)

USSR/Physics

Card 1/1 Pub. 22 - 14/49

Authors : Markov, M.A., Member-Correspondent of the Acad. of Sc., USSR

Title : "Hyperons", i.e., nucleons in an excited state, as a possible cause (mechanism) of mass particle formation

Periodical : Dok. AN SSSR 101/3, 449-452, Mar 21, 1955

Abstract : An attempt is made to connect (theoretically) the mass formation of particles (nucleons and anti-nucleons) with the disintegration of "hyperons", i.e., nucleons which are excited by π -mesons and are in the cascade form of disintegration. A number of peculiarities which may be met in the theories dealing with such subjects are pointed out (a few exemplary cases are discussed). Two references: 1 USSR and 1 USA (1949-1955).

Institution : The Acad. of Sc., USSR, P. N. Lebedev Physical Institute

Submitted : December 21, 1954

MARKOV, M.

Classification of hyperons and heavy mesons. M.
Markov, Soviet Phys. "Doklady" 1: 107-10 (1958) (Eng.
transl.)--See, C.A. 51, 870f. H. M. R.

proh
uji

mm Theory of Δ^0 particles: P. S. Isaev and M. A. Markov.
Soviet Phys. JETP 2, 84-8(1956) (Engl. translation) See
C.A. 49, 16535c. R.M.B.

2

Markov, M.
USSR/Theoretical Physics - Quantum Field Theory

B-6

Abst. Journal : Referat Zhur - Fizika, No 12, 1956, 33815

Author : Markov, M.

Institution : Physics Institute, Academy of Sciences USSR

Title : Remark Concerning the Systematization of Hyperons in Heavy Mesons

Original
Periodical : Dokl. AN SSSR, 1956, 106, No 5, 814-817

Abstract : A systematization of hyperons and K-mesons is proposed. The hyperons are considered as excited states of nucleons, whereby the number of the excitation coincides with the value of the oddity of the given hyperon. Generalizing the Fermi-Yang hypothesis, the author considers K-mesons as bound systems, consisting of a nucleon (anti-nucleon) and an antihyperon (hyperon). The suggested systematization is compared with the Gell-Mann systematization.

Card 1/1

MARKOV, M. A.

HYPERONS AND K MESONS. PART I. GENERAL CHAR-
ACTERISTICS OF HYPERONS AND K MESONS. M. A.

MARKOV, M. A. Institute of Nuclear Physics, Moscow State University, Moscow, U.S.S.R. 125851 and
1987. 102 p. 21 cm. 11551801 17

6
1-2 PM
1-8 PM
1-4E32

MARKOV, M. A.

19
HYPERONS AND K MESONS. PART II. ISOTOPIC SPIN
AND SYSTEMATICS OF FUNDAMENTAL PARTICLES. 19
M. A. Markov, Joint Institute of Nuclear Research. 116p.
and illustrations. Mar. 1967. (In Russian)
The isotopic invariants in various types of interactions,
the development of isotopic formalism and formal scheme
for classical particle classification, four dimensional vari-
ance of the isotopic spin, and further development of tri-
dimensional variance are discussed, as well as structural
concepts in the theory of fundamental particles and
M. Gell-Mann's scheme (Phys. Rev. 191, 433(1966)).
(R.V.J.)

6
1-1000
1-1000
4E3d

MARKOV, M

AUTHOR: MARKOV, M. 53-2-9/9
 TITLE: Comments to D.KONOSSENKO'S Work on "Semiconductor Bolometers"
 (Po povodu stat'i D.Konosenko: Poluprovodnikovyye bolometry, Russian)
 PERIODICAL: Uspekhi Fiz.Nauk, 1957, Vol 62, Nr 2, pp 197-199 (U.S.S.R.)

ABSTRACT: In his paper KONOSSENKO endeavors to systemize the bolometers made of semiconductors and to establish their importance for the measuring of infrared radiation as compared with other bolometers. According to MARKOV this statement contains a number of errors which he mentions and discusses:
 KONOSSENKO, when judging the sensitivity of bolometers, ignores the particular features of the electric amplifiers connected to the bolometer and therefore his evaluation of them is wrong. KONOSSENKO'S opinion that semiconductor bolometers are better than metal bolometers cannot be shared. MARKOV proves to KONOSSENKO that he did not display enough care when citing published works as references. Furthermore, MARKOV declares several of KONOSSENKO'S statements concerning various properties as absolutely wrong. KONOSSENKO'S paper appeared in Uspekhi Fizicheskikh Nauk 1955, Vol. 56, Nr 2, p.283.

PRESENTED BY:
 SUBMITTED:
 AVAILABLE: Library of Congress
 Card 1/1

21(8)

PHASE I BOOK EXPLOITATION

SOV/1413

Markov, Moisey Aleksandrovich

Giperony i K-mezony (Hyperons and K-Mesons) Moscow, Fizmatgiz, 1958.
343 p. (Series: Sovremennyye problemy fiziki) 5,000 copies printed.

Eds.: Alekseyev, D.M. and I.I. Denisov; Tech. Ed.: Akhlanov, S.N.

PURPOSE: This book is intended for theoretical physicists and scientists in the field of nuclear physics and the physics of elementary particles. It may also be used by students of advanced courses and Aspirants specializing in nuclear physics.

COVERAGE: A detailed review is given of experimental and theoretical work on the physics of two cosmic ray particles: hyperons and K-mesons.

TABLE OF CONTENTS

Foreword

PART I. GENERAL CHARACTERISTICS OF HYPERONS AND K-MESONS

7

Card 1/6

MARKOV, M. A.

"An Example of a Field Theory with Indefinite Metric in Hilbert Space."

Nuclear Physics, 10, 2, 1958, pp. 140-50 (No. Holland Publ. Co.,)

Joint Inst. of Nuclear Research, Dubna, USSR

A field theory is considered in which the Green's functions have singularities not on a cone but on a hyperboloid. The Metric of the corresponding Hilbert space is indefinite. Equations of motion for the boson and fermion fields are derived. The theory goes over to the usual field theory when a certain constant (characteristic length) approaches zero. The ultimate aim of the investigation is an attempt to find a possible reasonable interpretation of the negative energy density of the boson fields, of the negative probabilities, and of other similar features which appear in the theory, by appropriately modifying the concept of a particle coordinate so as to obtain the proposed propagation functions.